

Yemen*Wheat*

Standards	Seed Classes		
	Pre-basic	Basic	Certified
Field Standards			
Rotation (min. years)	2	1	1
Isolation (min. meters)			
Other varieties (max. %)	0.03	0.05	0.10
Other species (max. %)	0.01	0.03	0.05
Noxious weeds (max. %)	0.01	0.3	0.05
Infected plants (max. %)			
Seed Standards			
Pure seed (min. %)	98	97	96
Other crop seeds (max., %)	0.02	0.05	0.1
Weed seeds (max., %)	0.02	0.05	0.1
Inert matter (max. %)			
Infected seeds (max. %)			
Loose smut	None	0.05	0.1
Bunt	None	0.02	0.05
Germination (min. %)	85	85	85
Moisture content (max. %)	12	12	12

Yemen*Barley*

Standards	Seed Classes		
	Pre-basic	Basic	Certified
Field Standards			
Rotation (min. years)			
Isolation (min. meters)			
Other varieties (max. %)	0.03	0.05	0.10
Other species (max. %)	0.01	0.03	0.05
Noxious weeds (max. %)	0.01	0.3	0.05
Infected plants (max. %)			
Seed Standards			
Pure seed (min. %)	98	97	96
Other crop seeds (max. %)	0.02	0.05	0.1
Weed seeds (max. %)	0.02	0.05	0.1
Inert matter (max. %)			
Infected seeds (max. %)			
Loose smut	None	0.05	0.1
Bunt	None	0.02	0.05
Germination (min. %)	85	85	85
Moisture content (max. %)	12	12	12

Yemen*Maize*

Standards	Seed Classes		
	Pre-basic	Basic	Certified
Field Standards			
Rotation (min. years)			
Isolation (min. meters)			
Other varieties (max. %)	None	1	2
Other species (max. %)		0.02	0.5
Noxious weeds (max. %)			
Infected plants (max. %)			
Seed Standards			
Pure seed (min. %)		98	97
Other crop seeds (max. %)			0.05
Weed seeds (max. %)			
Inert matter (max. %)			
Infected seeds (max. %)			
Smut	0.5	0.1	0.2
Germination (min. %)		85	85
Moisture content (max. %)	13	13	13

Yemen*Sorghum*

Standards	Seed Classes		
	Pre-basic	Basic	Certified
Field Standards			
Rotation (min. years)			
Isolation (min. meters)			
Other varieties (max. %)		0.02	0.05
Other species (max. %)		0.01	0.02
Noxious weeds (max. %)			
Infected plants (max. %)			
Seed Standards			
Pure seed (min. %)		95	93
Other crop seeds (max. %)		0.1	0.2
Weed seeds (max. %)		0.1	0.2
Inert matter (max. %)			
Infected seeds (max. %)			
Smut	0.5	0.1	0.2
Germination (min. %)		80	80
Moisture content (max. %)		13	13

Yemen*Millet*

Standards	Seed Classes		
	Pre-basic	Basic	Certified
Field Standards			
Rotation (min. years)			
Isolation (min. meters)			
Other varieties (max. %)		0.02	0.05
Other species (max. %)		0.01	0.02
Noxious weeds (max. %)			
Infected plants (max. %)			
Seed Standards			
Pure seed (min. %)		95	93
Other crop seeds (max., %)		0.1	0.2
Weed seeds (max., %)		0.1	0.2
Inert matter (max. %)			
Infected seedss (max., %)			
Germination (min. %)		80	80
Moisture content (max. %)		13	13

Yemen

Potato

Standards	Seed Classes		
	Basic seed (E)	Certified (A)	Certified (B)

Field Standards

- Rotation (min. years)
- Isolation (min. meters)
- Other Varieties (max. %)
- Other species (max. %)
- Noxious weeds (max. %)
- Infected plants (max. %)

Seed Standards (tubers)

Dry rot caused by Alternaria, Fusarium Phoma Phytophtra spp	1	2	3
Wet rots Black leg	None	1	1
Rhizoctonia Heavy infection Total infection			
Insects Tuber moth Heavy Medium Light	None None None	0.5 2 5	.05 2 5
Cut worms	1	5	5
